

宁夏饲料营养数据库

表 1 饲料描述及常规成分

| 序号 | 饲料号 | 饲料名称 | 饲料描述 | 干物质 DM% | 粗蛋白质 CP% | 粗脂肪 EE% | 粗纤维 CF% | 无氮浸出物 NFE% | 粗灰分 Ash% | 中洗纤维 NDF% | 酸洗纤维 ADF% | 淀粉 Starch% | 钙 Ca% | 总磷 P% | 有效磷 P% |
|----|-----------|-------|------------------------------|---------|----------|---------|---------|------------|----------|-----------|-----------|------------|-------|-------|--------|
| 1 | 4-07-0278 | 玉米 | 成熟,高蛋白,优质 | 88.0 | 9.0 | 3.5 | 2.8 | 71.5 | 1.2 | 9.1 | 3.3 | 61.7 | 0.01 | 0.31 | 0.09 |
| 2 | 4-07-0288 | 玉米 | 成熟,高赖氨酸,优质 | 86.0 | 8.5 | 5.3 | 2.6 | 68.3 | 1.3 | 9.4 | 3.5 | 59.0 | 0.16 | 0.25 | 0.05 |
| 3 | 4-07-0279 | 玉米 | 成熟,GB1353-2018,1 级 | 86.0 | 8.7 | 3.6 | 1.6 | 70.7 | 1.4 | 9.3 | 2.7 | 65.4 | 0.02 | 0.27 | 0.05 |
| 4 | 4-07-0280 | 玉米 | 成熟,GB1353-2018,2 级 | 86.0 | 8.0 | 3.6 | 2.3 | 71.8 | 1.2 | 9.9 | 3.1 | 63.5 | 0.02 | 0.27 | 0.05 |
| 5 | 4-07-0272 | 高粱 | 成熟,GB 8231-87 | 88.0 | 8.7 | 3.4 | 1.4 | 70.7 | 1.8 | 17.4 | 8.0 | 68.0 | 0.13 | 0.36 | 0.09 |
| 6 | 4-07-0270 | 小麦 | 混合小麦,成熟 GB1351-2008 2 级 | 88.0 | 13.4 | 1.7 | 1.9 | 69.1 | 1.9 | 13.3 | 3.9 | 54.6 | 0.17 | 0.41 | 0.21 |
| 7 | 4-07-0274 | 大麦(裸) | 裸大麦,成熟 GB/T 11760-2008 2 级 | 87.0 | 13.0 | 2.1 | 2.0 | 67.7 | 2.2 | 10.0 | 2.2 | 50.2 | 0.04 | 0.39 | 0.12 |
| 8 | 4-07-0277 | 大麦(皮) | 皮大麦,成熟 GB 10367-89 1 级 | 87.0 | 11.0 | 1.7 | 4.8 | 67.1 | 2.4 | 18.4 | 6.8 | 52.2 | 0.09 | 0.33 | 0.10 |
| 9 | 4-07-0281 | 黑麦 | 籽粒,进口 | 88.0 | 9.5 | 1.5 | 2.2 | 73.0 | 1.8 | 12.3 | 4.6 | 56.5 | 0.05 | 0.30 | 0.14 |
| 10 | 4-07-0273 | 稻谷 | 成熟,晒干 NY/T 2 级 | 86.0 | 7.8 | 1.6 | 8.2 | 63.8 | 4.6 | 27.4 | 13.7 | 63.0 | 0.03 | 0.36 | 0.15 |
| 11 | 4-07-0276 | 糙米 | 除去外壳的大米, GB/T 18810-2002,1 级 | 87.0 | 8.8 | 2.0 | 0.7 | 74.2 | 1.3 | 1.6 | 0.8 | 47.8 | 0.03 | 0.35 | 0.13 |
| 12 | 4-07-0275 | 碎米 | 加工精米后的副产品 GB/T 5503-2009 1 级 | 88.0 | 10.4 | 2.2 | 1.1 | 72.7 | 1.6 | 0.8 | 0.6 | 51.6 | 0.06 | 0.35 | 0.12 |
| 13 | 4-07-0479 | 粟(谷子) | 合格,带壳,成熟 | 86.5 | 9.7 | 2.3 | 6.8 | 65.0 | 2.7 | 15.2 | 13.3 | 63.2 | 0.12 | 0.30 | 0.09 |
| 14 | 4-04-0067 | 木薯干 | 木薯干片,晒干 GB 10369-89 合格 | 87.0 | 2.5 | 0.7 | 2.5 | 79.4 | 1.9 | 8.4 | 6.4 | 71.6 | 0.27 | 0.09 | 0.03 |
| 15 | 4-04-0068 | 甘薯干 | 甘薯干片,晒干 NY/T 121-1989 合格 | 87.0 | 4.0 | 0.8 | 2.8 | 76.4 | 3.0 | 8.1 | 4.1 | 64.5 | 0.19 | 0.02 | — |
| 16 | 4-08-0104 | 次粉 | 黑面,黄粉,下面 NY/T 211-1992 1 级 | 88.0 | 15.4 | 2.2 | 1.5 | 67.1 | 1.5 | 18.7 | 4.3 | 37.8 | 0.08 | 0.48 | 0.17 |
| 17 | 4-08-0105 | 次粉 | 黑面,黄粉,下面 NY/T 211-1992 2 级 | 87.0 | 13.6 | 2.1 | 2.8 | 66.7 | 1.8 | 31.9 | 10.5 | 36.7 | 0.08 | 0.48 | 0.17 |
| 18 | 4-08-0069 | 小麦麸 | 传统制粉工艺 GB 10368-1989 1 级 | 87.0 | 15.7 | 3.9 | 6.5 | 56.0 | 4.9 | 37.0 | 13.0 | 22.6 | 0.11 | 0.92 | 0.32 |
| 19 | 4-08-0070 | 小麦麸 | 传统制粉工艺 GB 10368-1989 2 级 | 87.0 | 14.3 | 4.0 | 6.8 | 57.1 | 4.8 | 41.3 | 11.9 | 19.8 | 0.10 | 0.93 | 0.33 |
| 20 | 4-08-0041 | 米糠 | 新鲜,不脱脂 NY/T 2 级 | 90.0 | 14.5 | 15.5 | 6.8 | 45.6 | 7.6 | 20.3 | 11.6 | 27.4 | 0.05 | 2.37 | 0.35 |
| 21 | 4-10-0025 | 米糠饼 | 未脱脂,机榨 NY/T 1 级 | 90.0 | 15.0 | 9.2 | 7.6 | 49.3 | 8.9 | 28.3 | 11.9 | 30.9 | 0.14 | 1.73 | 0.25 |
| 22 | 4-10-0018 | 米糠粕 | 浸提或预压浸提, NY/T 1 级 | 87.0 | 15.1 | 2.0 | 7.5 | 53.6 | 8.8 | 23.3 | 10.9 | 25.0 | 0.15 | 1.82 | 0.25 |
| 23 | 5-09-0127 | 大豆 | 黄大豆,成熟 GB 1352-86 2 级 | 87.0 | 35.5 | 17.3 | 4.3 | 25.7 | 4.2 | 7.9 | 7.3 | 2.6 | 0.27 | 0.48 | 0.12 |

续表 1

| 序号 | 饲料号 | 饲料名称 | 饲料描述 | 干物质 DM% | 粗蛋白质 CP% | 粗脂肪 EE% | 粗纤维 CF% | 无氮浸出物 NFE% | 粗灰分 Ash% | 中洗纤维 NDF% | 酸洗纤维 ADF% | 淀粉 Starch% | 钙 Ca% | 总磷 P% | 有效磷 A-P% |
|----|-----------|-------|---------------------------------------|------------|-------------|------------|------------|---------------|-------------|--------------|--------------|---------------|----------|----------|-------------|
| 24 | 5-09-0128 | 全脂大豆 | 微粒化 GB/T 20411-2006 | 88.0 | 35.5 | 18.7 | 4.6 | 25.2 | 4.0 | 11.0 | 6.4 | 6.7 | 0.32 | 0.40 | 0.10 |
| 25 | 5-10-0241 | 大豆饼 | 机榨 GB 10379-`989 2 级 | 89.0 | 41.8 | 5.8 | 4.8 | 30.7 | 5.9 | 18.1 | 15.5 | 3.6 | 0.31 | 0.50 | 0.13 |
| 26 | 5-10-0103 | 去皮大豆粕 | 去皮,浸提或预压浸提NY/T 1 级 | 89.0 | 47.9 | 1.5 | 3.3 | 29.7 | 4.9 | 8.8 | 5.3 | 1.8 | 0.34 | 0.65 | 0.24 |
| 27 | 5-10-0102 | 大豆粕 | 浸提或预压浸提GB/T19541-2017 | 89.0 | 44.2 | 1.9 | 5.9 | 28.3 | 6.1 | 13.6 | 9.6 | 3.5 | 0.33 | 0.62 | 0.16 |
| 28 | 5-10-0118 | 棉籽饼 | 机榨NY/ T 129-1989 2 级 | 88.0 | 36.3 | 7.4 | 12.5 | 26.1 | 5.7 | 32.1 | 22.9 | 3.0 | 0.21 | 0.83 | 0.21 |
| 29 | 5-10-0119 | 棉籽粕 | 浸提 GB 21264-2007 1 级 | 90.0 | 47.0 | 0.5 | 10.2 | 26.3 | 6.0 | 22.5 | 15.3 | 1.5 | 0.25 | 1.10 | 0.28 |
| 30 | 5-10-0117 | 棉籽粕 | 浸提 GB 21264-2007 2 级 | 90.0 | 43.5 | 0.5 | 10.5 | 28.9 | 6.6 | 28.4 | 19.4 | 1.8 | 0.28 | 1.04 | 0.26 |
| 31 | 5-10-0220 | 棉籽蛋白 | 脱酚,低温一次浸出,分步萃取 | 92.0 | 51.1 | 1.0 | 6.9 | 27.3 | 5.7 | 20.0 | 13.7 | 0.9 | 0.29 | 0.89 | 0.22 |
| 32 | 5-10-0183 | 菜籽饼 | 机榨NY/ T 1799-2009 2 级 | 88.0 | 35.7 | 7.4 | 11.4 | 26.3 | 7.2 | 33.3 | 26.0 | 3.8 | 0.59 | 0.96 | 0.20 |
| 33 | 5-10-0121 | 菜籽粕 | 浸提GB/T 23736-2009 2 级 | 88.0 | 38.6 | 1.4 | 11.8 | 28.9 | 7.3 | 20.7 | 16.8 | 6.1 | 0.65 | 1.02 | 0.25 |
| 34 | 5-10-0116 | 花生仁饼 | 机榨NY/ T 132-2019 2 级 | 88.0 | 44.7 | 7.2 | 5.9 | 25.1 | 5.1 | 14.0 | 8.7 | 6.6 | 0.25 | 0.53 | 0.16 |
| 35 | 5-10-0115 | 花生仁粕 | 浸提GB 10382-1989 2 级 | 88.0 | 47.8 | 1.4 | 6.2 | 27.2 | 5.4 | 15.5 | 11.7 | 6.7 | 0.27 | 0.56 | 0.17 |
| 36 | 1-10-0031 | 向日葵仁饼 | 壳仁比 35:65 NY/ T 128-1989 3 级 | 88.0 | 29.0 | 2.9 | 20.4 | 31.0 | 4.7 | 41.4 | 29.6 | 2.0 | 0.24 | 0.87 | 0.22 |
| 37 | 5-10-0242 | 向日葵仁粕 | 壳仁比 16:84 GB 10377-89 T 2 级 | 88.0 | 36.5 | 1.0 | 10.5 | 34.4 | 5.6 | 14.9 | 13.6 | 6.2 | 0.27 | 1.13 | 0.29 |
| 38 | 5-10-0243 | 向日葵仁粕 | 壳仁比 24:76 GB 10377-89 T 2 级 | 88.0 | 33.6 | 1.0 | 14.8 | 38.8 | 5.3 | 32.8 | 23.5 | 4.4 | 0.26 | 1.03 | 0.26 |
| 39 | 5-10-0119 | 亚麻仁饼 | 机榨NY/ T 216-1992 2 级 | 88.0 | 32.2 | 7.8 | 7.8 | 34.0 | 6.2 | 29.7 | 27.1 | 11.4 | 0.39 | 0.88 | 0.22 |
| 40 | 5-10-0120 | 亚麻仁粕 | 浸提或 预压 浸提 NY/T 215-1992 2 级 | 88.0 | 34.8 | 1.8 | 8.2 | 36.6 | 6.6 | 21.6 | 14.4 | 13.0 | 0.42 | 0.95 | 0.24 |
| 41 | 5-10-0246 | 芝麻饼 | 机榨,CP40% | 92.0 | 39.2 | 10.3 | 7.2 | 24.9 | 10.4 | 18.0 | 13.2 | 1.8 | 2.24 | 1.19 | 0.31 |
| 42 | 5-11-0001 | 玉米蛋白粉 | 去胚芽、淀粉后的面筋部分 NY 685-2003 1 级 | 90.1 | 63.5 | 5.4 | 1.0 | 19.2 | 1.0 | 8.7 | 4.6 | 17.2 | 0.07 | 0.44 | 0.16 |
| 43 | 5-11-0002 | 玉米蛋白粉 | 同上,中等蛋白质产品,CP50% NY/T 685-2003 2 级 | 88.0 | 56.3 | 4.7 | 1.3 | 23.4 | 2.3 | 8.2 | 5.1 | 16.1 | 0.04 | 0.44 | 0.15 |
| 44 | 5-11-0008 | 玉米蛋白粉 | 同上,中等蛋白质产品,CP40% NY/T 685-2003 3 级 | 89.9 | 44.3 | 6.0 | 1.6 | 37.1 | 0.9 | 29.1 | 8.2 | 20.6 | 0.12 | 0.50 | 0.31 |

| 序号 | 饲料号 | 饲料名称 | 饲料描述 | 干物质 DM% | 粗蛋白 质 CP% | 粗脂肪 EE% | 粗纤维 CF% | 无氮浸出物 NFE% | 粗灰分 Ash% | 中洗纤维 NDF% | 酸洗纤维 ADF% | 淀粉 Starch% | 钙 Ca% | 总磷 P% | 有效磷 A-P% |
|----|-----------|-------------|--------------------------|------------|-----------------|------------|------------|---------------|-------------|--------------|--------------|---------------|----------|----------|-------------|
| 45 | 5-11-0003 | 玉米蛋白饲料 | 玉米去胚芽、淀粉后的含皮残渣 | 88.0 | 18.3 | 7.5 | 7.8 | 47.0 | 5.4 | 33.6 | 10.5 | 21.5 | 0.15 | 0.70 | 0.17 |
| 46 | 4-10-0026 | 玉米胚芽饼 | 玉米湿磨后的胚芽,机榨 | 90.0 | 16.7 | 9.6 | 6.3 | 50.8 | 6.6 | 28.5 | 7.4 | 13.5 | 0.04 | 0.50 | 0.15 |
| 47 | 4-10-0244 | 玉米胚芽粕 | 玉米湿磨后的胚芽,浸提 | 90.0 | 20.8 | 2.0 | 6.5 | 54.8 | 5.9 | 38.2 | 10.7 | 14.2 | 0.06 | 0.50 | 0.15 |
| 48 | 5-11-0007 | DDGS | 玉米酒精糟及可溶物,脱水 | 89.2 | 27.5 | 10.1 | 6.6 | 39.9 | 5.1 | 38.3 | 12.5 | 4.2 | 0.06 | 0.71 | 0.48 |
| 49 | 5-11-0009 | 蚕豆粉浆蛋白粉 | 蚕豆去皮制粉丝后的浆液,脱水 | 88.0 | 66.3 | 4.7 | 4.1 | 10.3 | 2.6 | 13.7 | 9.7 | - | | 0.59 | 0.18 |
| 50 | 5-11-0004 | 麦芽根 | 大麦芽副产品,干燥 | 89.7 | 28.3 | 1.4 | 12.5 | 41.4 | 6.1 | 40.0 | 15.1 | 7.2 | 0.22 | 0.73 | 0.18 |
| 51 | 5-13-0044 | 鱼粉(CP67%) | 进口GB/T 19164-2003,特级 | 92.4 | 67.0 | 8.4 | 0.2 | 0.4 | 16.4 | | | | 4.56 | 2.88 | 2.88 |
| 52 | 5-13-0046 | 鱼粉(CP60.2%) | 沿海产的海鱼粉,脱脂12 样平均值 | 90.0 | 60.2 | 4.9 | 0.5 | 11.6 | 12.8 | | | | 4.04 | 2.90 | 2.90 |
| 53 | 5-13-0077 | 鱼粉(CP53.5%) | 沿海产的海鱼粉,脱脂11 样平均值 | 90.0 | 53.5 | 10.0 | 0.8 | 4.9 | 20.8 | | | | 5.88 | 3.20 | 3.20 |
| 54 | 5-13-0036 | 血粉 | 鲜猪血 喷雾干燥, 国产 | 88.0 | 82.8 | 0.4 | | 1.6 | 3.2 | | | | 0.29 | 0.31 | 0.29 |
| 55 | 5-13-0037 | 羽毛粉 | 纯净羽毛,水解, 国产 | 88.0 | 77.9 | 2.2 | 0.7 | 1.4 | 5.8 | | | | 0.20 | 0.68 | 0.61 |
| 56 | 5-13-0038 | 皮革粉 | 废牛皮,水解, 国产 | 88.0 | 74.7 | 0.8 | 1.6 | | 10.9 | | | | 4.40 | 0.15 | 0.13 |
| 57 | 5-13-0047 | 肉骨粉 | 屠宰下脚,带骨干燥粉碎 | 93.0 | 50.0 | 8.5 | 2.8 | | 31.7 | | | | 9.20 | 4.70 | 4.37 |
| 58 | 5-13-0048 | 肉粉 | 脱脂, 国产 | 94.0 | 54.0 | 12.0 | 1.4 | 4.3 | 22.3 | | | | 7.69 | 3.88 | 3.61 |
| 59 | 1-05-0074 | 苜蓿草粉(CP19%) | 一茬盛花期烘干 NY/T 140-2002 1级 | 87.0 | 19.1 | 2.3 | 22.7 | 35.3 | 7.6 | 36.7 | 25.0 | 6.1 | 1.40 | 0.51 | 0.51 |
| 60 | 1-05-0075 | 苜蓿草粉(CP17%) | 一茬盛花期烘干 NY/T 140-2002 2级 | 87.0 | 17.2 | 2.6 | 25.6 | 33.3 | 8.3 | 39.0 | 28.6 | 3.4 | 1.52 | 0.22 | 0.22 |
| 61 | 1-05-0076 | 苜蓿草粉 | NY/T 140-2002 3 级 | 87.0 | 14.3 | 2.1 | 29.8 | 33.8 | 10.1 | 36.8 | 29.0 | 3.5 | 1.34 | 0.19 | 0.19 |
| 62 | 5-11-0005 | 啤酒糟 | 大麦酿造副产品 | 88.0 | 24.3 | 5.3 | 13.4 | 40.8 | 4.2 | 39.4 | 24.6 | 11.5 | 0.32 | 0.42 | 0.14 |
| 63 | 7-15-0001 | 啤酒酵母 | 啤酒酵母菌粉,QB/T1940-94 | 91.7 | 52.4 | 0.4 | 0.6 | 33.6 | 4.7 | 6.1 | 1.8 | 1.0 | 0.16 | 1.02 | 0.46 |
| 64 | 4-13-0075 | 乳清粉 | 乳清,脱水,乳糖含量 73% | 97.2 | 11.5 | 0.8 | 0.1 | 76.8 | 8.0 | | | | 0.62 | 0.69 | 0.52 |
| 65 | 5-01-0162 | 酪蛋白 | 脱水, 来源于牛奶 | 91.7 | 89.0 | 0.2 | | 0.4 | 2.1 | | | | 0.20 | 0.68 | 0.67 |
| 66 | 5-14-0503 | 明胶 | 食用 | 90.0 | 88.6 | 0.5 | | 0.59 | 0.31 | | | | 0.49 | | |
| 67 | 4-06-0076 | 牛奶乳糖 | 进口,含乳糖 80%以上 | 96.0 | 3.5 | 0.5 | | 82.0 | 10.0 | | | | 0.52 | 0.62 | 0.62 |
| 68 | 4-06-0077 | 乳糖 | 食用 | 96.0 | 0.3 | | | 95.7 | | | | | | | |
| 69 | 4-06-0078 | 葡萄糖 | 食用 | 90.0 | 0.3 | | | 89.7 | | | | | | | |

| 序号 | 饲料号 | 饲料名称 | 饲料描述 | 干物质 DM% | 粗蛋白质 CP% | 粗脂肪 EE% | 粗纤维 CF% | 无氮浸出物 NFE% | 粗灰分 Ash% | 中洗纤维 NDF% | 酸洗纤维 ADF% | 淀粉 Starch% | 钙 Ca% | 总磷 P% | 有效磷 A-P% |
|----|-----------|------|------|------------|-------------|------------|------------|---------------|-------------|--------------|--------------|---------------|----------|----------|-------------|
| 70 | 4-06-0079 | 蔗糖 | 食用 | 99.0 | | | | 98.5 | 0.5 | | | | 0.04 | | |
| 71 | 4-02-0889 | 玉米淀粉 | 食用 | 99.0 | 0.3 | 0.2 | | 98.5 | | | | 98.0 | | 0.03 | 0.01 |
| 72 | 4-17-0001 | 牛脂 | | 99.0 | | 98.0* | | 0.5 | 0.5 | | | | | | |
| 73 | 4-17-0002 | 猪油 | | 99.0 | | 98.0* | | 0.5 | 0.5 | | | | | | |
| 74 | 4-17-0003 | 家禽脂肪 | | 99.0 | | 98.0* | | 0.5 | 0.5 | | | | | | |
| 75 | 4-17-0004 | 鱼油 | | 99.0 | | 98.0* | | 0.5 | 0.5 | | | | | | |
| 76 | 4-17-0005 | 菜籽油 | | 99.0 | | 98.0* | | 0.5 | 0.5 | | | | | | |
| 77 | 4-17-0006 | 椰子油 | | 99.0 | | 98.0* | | 0.5 | 0.5 | | | | | | |
| 78 | 4-07-0007 | 玉米油 | | 99.0 | | 98.0* | | 0.5 | 0.5 | | | | | | |
| 79 | 4-17-0008 | 棉籽油 | | 99.0 | | 98.0* | | 0.5 | 0.5 | | | | | | |
| 80 | 4-17-0009 | 棕榈油 | | 99.0 | | 98.0* | | 0.5 | 0.5 | | | | | | |
| 81 | 4-17-0010 | 花生油 | | 99.0 | | 98.0* | | 0.5 | 0.5 | | | | | | |
| 82 | 4-17-0011 | 芝麻油 | | 99.0 | | 98.0* | | 0.5 | 0.5 | | | | | | |
| 83 | 4-17-0012 | 大豆油 | 粗制 | 99.0 | | 98.0* | | 0.5 | 0.5 | | | | | | |
| 84 | 4-17-0013 | 葵花油 | | 99.0 | | 98.0* | | 0.5 | 0.5 | | | | | | |

续表 1

* ①代表典型值（下同）；②“*”空的数据项代表为“0”（下同）；③从表 1~表 12 所示所有数据，无特别说明者，均表示为饲喂状态的含量数据。